

Participant ID: Participant Initials:

Clinical Center: Visit Number: Site:

CRF Date: RC ID:

	SCREENING LABORATORY RESULTS					
No	Note: Medical record is defined as a clinical value collected no more than 90 days prior to screening visit.					
Blood Collection:						
1.	Source of serum creatinine result:	□₁ Medical Records□₂ Blood drawn at screening visit				
2.	Date of Result/Blood draw:	/ / (mm/dd/yyyy)				
3.	Serum creatinine value (Site-based laboratory results):	mg/dL				
Skip to Question #8 if source of serum creatinine is from "Medical Records". Continue to Questions #4 if source of serum creatinine is "Blood drawn at screening visit".						
4.	Time of blood draw:	: (military time)				
5.	Institution Laboratory Code:					
6.	Is this a non-fasting blood sample?	\square_1 Yes \square_0 No				
7.	Blood sample reserved for Central Laboratory?	□ ₁ Yes □ ₀ No				
Ur	ine Collection:					
8.	Date of spot urine test at screening visit.	/ / (mm/dd/yyyy)				
9.	Results of spot urine test at screening visit.					
	a. Glucose:	□ ₁ Positive □ ₀ Negative □ ₉₇ Not Done				
	b. Protein:	\square_0 Negative \square_1 Trace \square_2 1+ \square_3 2+ \square_4 3+ \square_5 4+				
	c. Hematuria:	□₁ Positive □₀ Negative □₃γ Not Done				

SCRLABII Page 1 of 2 V3.0.20140616

R	ENAL	NSUA	
N. C.	\subseteq		à
FRO C		ic	ENC
O	<u> </u>		7
O	/0-	TUD.	

Participant ID: Participant Initials:

Clinical Center: Site: Visit Number:

CRF Date: RC ID:

SCREENING LABORATORY RESULTS

Proteinuria Data From Medical Records:

Only proteinuria data from the medical record obtained within 90 days of the screening visit may be recorded.

10. Is there documentation of sufficient proteinuria (see table below) in the participant's medical record within the past 90 days?	□ ₁ Yes	□ ₀ No	□ ₉₉ No documentation obtained
a. If Yes , date of proteinuria result from medical record:	/	/	(mm/dd/yyyy)

Measure:	Threshold Defining Sufficient Proteinuria:
Albumin excretion rate (AER):	>300 mg/24 hours
Protein excretion rate (PER):	>500 mg/24 hours
Albumin-to-creatinine ratio (ACR):	>30 mg/mmol >300 mg/g
Protein-to-creatinine ratio (PCR):	>50 mg/mmol >500 mg/g
Protein reagent strip ('dipstick'):	1+ or greater

V3.0.20140616 Page 2 of 2 **SCRLABII**