

The Maryland Genetics of Interstitial Cystitis Study
The MaGIC study

Categories of Interstitial Cystitis/Painful Bladder Syndrome *

Definite IC/PBS

- 1) Female AND
- 2) Bladder pain AND
- 3) ≥ 2 pertinent urinary symptoms AND
- 4) Symptoms in 2) and 3) occur together and on most days for at least 4 weeks AND
- 5) No NIDDK exclusionary diseases, or neurogenic bladder

Probable

- 1) As Definite but only one pertinent urinary symptom OR
- 2) As Definite but non-bladder pelvic pain, not associated with GI symptoms OR
- 3) As Definite but Male

Possible

- 1) As Definite but no pertinent urinary symptoms OR
- 2) As Definite but non-bladder pelvic pain, not associated with GI symptoms AND one pertinent urinary symptom OR
- 3) As Definite but Male with bladder pain alone or with one pertinent urinary symptom OR
- 4) As Definite but Male with non-bladder pelvic pain, not associated with GI symptoms and ≥ 1 pertinent urinary symptoms

Uncertain

Male or Female with pelvic pain alone ≥ 4 weeks

Not IC/PBS

No pelvic pain ≥ 4 weeks

***Participants will be asked to describe these symptoms in the absence of pregnancy, high fluid intake, diuretic administration, urinary tract infection, or another known disease causing these symptoms**