

The Maryland Genetics of Interstitial Cystitis Study  
The MaGIC study

Categories of Interstitial Cystitis/Painful Bladder Syndrome \*

**Definite IC/PBS**

- 1) Female AND
- 2) Bladder pain AND
- 3)  $\geq 2$  pertinent urinary symptoms AND
- 4) Symptoms in 2) and 3) occur together and on most days for at least 4 weeks AND
- 5) No NIDDK exclusionary diseases, or neurogenic bladder

**Probable**

- 1) As Definite but only one pertinent urinary symptom OR
- 2) As Definite but non-bladder pelvic pain, not associated with GI symptoms OR
- 3) As Definite but Male

**Possible**

- 1) As Definite but no pertinent urinary symptoms OR
- 2) As Definite but non-bladder pelvic pain, not associated with GI symptoms AND one pertinent urinary symptom OR
- 3) As Definite but Male with bladder pain alone or with one pertinent urinary symptom OR
- 4) As Definite but Male with non-bladder pelvic pain, not associated with GI symptoms and  $\geq 1$  pertinent urinary symptoms

**Uncertain**

Male or Female with pelvic pain alone  $\geq 4$  weeks

**Not IC/PBS**

No pelvic pain  $\geq 4$  weeks

**\*Participants will be asked to describe these symptoms in the absence of pregnancy, high fluid intake, diuretic administration, urinary tract infection, or another known disease causing these symptoms**