



Participant ID: _____

Pin # _____

Discovery Site: _____

Clinical Center _____

CRF Date: ____/____/____

Visit #: _____

COMPLEX MEDICAL SYMPTOMS INVENTORY – TMJ

MODULE	DIAGNOSIS CRITERIA CITATION	SYMPTOM QUESTION	SUPPORTING SYMPTOMS/ QUALIFIERS	PHYSICAL EXAM	NOTES
M: TMJ	Temporomandibular Joint Disorder Dworkin, SF (1992) Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. J Craniomandib Disord Facial Oral pain. 6(4):327-355.	(Sx 31) Jaw and/or face pain (Sx 32) Temple pain (Sx 23) Ear pain	Tight jaw	Tenderpoint exam • ≥3 TP to 1kg of digital pressure • ≥1 TP must be ipsilateral to complaint of pain	<ul style="list-style-type: none"> • Must have at least 1/3 symptoms in addition to a positive TP exam Myofacial TMDJ with limited opening: fulfilling above criteria, plus Pain-free unassisted mandibular opening of <40 mm, plus Maximum assisted opening (passive stretch) of ≥5 mm greater than pain-free unassisted opening

<input type="checkbox"/> ₁ Yes	<input type="checkbox"/> ₀ No	7A	Jaw and/or face pain (Sx 31)	If YES to 7A, 7B, or 7C, perform tenderpoint exam.
<input type="checkbox"/> ₁ Yes	<input type="checkbox"/> ₀ No	7B	Temple pain (Sx 32)	
<input type="checkbox"/> ₁ Yes	<input type="checkbox"/> ₀ No	7C	Ear pain (Sx 23)	
<input type="checkbox"/> ₁ Yes	<input type="checkbox"/> ₀ No	7D	>= 3 tenderpoints	
<input type="checkbox"/> ₁ Yes	<input type="checkbox"/> ₀ No	M:TMJ	Diagnosis: YES to ≥1/3 7A-7C and 7D	
<input type="checkbox"/> ₁ Yes	<input type="checkbox"/> ₀ No	7E	TMJD with limited ROM = YES CMSI-TMJ and pain-free unassisted mandibular opening of <40mm, plus maximum assisted opening (passive stretch) of >=5mm greater than pain-free unassisted opening	
<input type="checkbox"/> ₁ Yes	<input type="checkbox"/> ₀ No	7F	TMJD without limited ROM = YES to CMSI-TMJ	