|                | _ |  |  |      |      |    |  |  |  |
|----------------|---|--|--|------|------|----|--|--|--|
| Participant ID |   |  |  | Nicl | knam | ne |  |  |  |

RISE **OGIT.1**July 2013
Page 1 of 3



## Restoring Insulin Secretion Study OGTT: Oral Glucose Tolerance Test

| 1.  | Study Visit Number VISIT  | BAS M06 M12 M15 M21                |  |  |  |  |
|---|---|------------------------------------|--|--|--|--|
| 2.  | Visit date (mm/dd/yyyy) Replaced with DAYSRAND  |                                    |  |  |  |  |
| 3.  | Staff ID  |                                    |  |  |  |  |
|   |   |                                    |  |  |  |  |
| Ins   | tructions: Complete this form at any visit when an OGTT is co   | ompleted (BAS, M06, M12, M15, M21) |  |  |  |  |
| Tes   | t Qualification   |                                    |  |  |  |  |
| Со  | nfirm the participant's readiness for the OGTT:   |                                    |  |  |  |  |
|   | a. Did not take any study medications the morning of this visit   | s True False                       |  |  |  |  |
|   | b. Had no illness within past week  | True False                         |  |  |  |  |
|   | c. Did not exercise, other than walk, within past 10 hours  | True False                         |  |  |  |  |
|   | d. Has fasted at least 10 hours   | True False                         |  |  |  |  |
|   | e. Had regular diet over last 3 days  | True False                         |  |  |  |  |
| 5.  | Is the participant prepared for the oral glucose tolerance test? <i>All above must be true</i> . OGPREP | Yes 2 No                           |  |  |  |  |
| 6.  | Fasting finger-stick blood sugar elevated (see MOP v.2 4.4 for details)? OGFAST                         | 4 1 Yes 2 No                       |  |  |  |  |
|   | → If YES, see MOP Volume 2 Section 4.4 for information or continuing or rescheduling visit.             | n                                  |  |  |  |  |
| Tes   | t Progression   |                                    |  |  |  |  |
| 7.  | Time of 1st fasting blood sample OGFSTTIME1 (-10 min)   | : 24 hour clock                    |  |  |  |  |
| 8.  | Time of 2 <sup>nd</sup> fasting blood sample OGFSTTIME2 (-5 min)  | : 24 hour clock                    |  |  |  |  |
|   | OGIT Samples  |                                    |  |  |  |  |
|   | Time glucose consumption started  → Start timer now   | 0 0 0 : 0 0 Timer                  |  |  |  |  |
| 9.  | Time glucose consumption ended OGGLEND1 / OGGLUEND2   | : Timer                            |  |  |  |  |
| → Glucola should be completely consumed within 5 minutes. |   |                                    |  |  |  |  |

| Participant ID Nickname   | RISE <b>OGTT.1</b> July 2013 Page 2 of 3 |  |  |  |  |  |  |
|---|--|--|--|--|--|--|--|
| Test Progression (continued)  |  |  |  |  |  |  |  |
| The following samples should be drawn within 2 minutes of intervals (before or after) counting from the time the glucose drink was started. If a sample is drawn outside that window, record the time and continue with the OGTT. |  |  |  |  |  |  |  |
| 10. Time 10 minute sample drawn OGDRAW10M / OGDRAW10S   | : Timer                                  |  |  |  |  |  |  |
| 11. Time 20 minute sample drawn OGDRAW20M/OGDRAW20S   | : Timer                                  |  |  |  |  |  |  |
| 12. Time 30 minute sample drawn OGDRAW30M/OGDRAW30S   | : Timer                                  |  |  |  |  |  |  |
| 13. Time 60 minute sample drawn OGDRAW60M / OGDRAW60S (1 hour)  | : Timer                                  |  |  |  |  |  |  |
| 14. Time 90 minute sample drawn OGDRAW90M / OGDRAW90S   | : Timer                                  |  |  |  |  |  |  |
| 15. Time 120 minute sample drawn OGDRAW120M / OGDRAW120S (2 hour)   | Timer                                    |  |  |  |  |  |  |
| 16. Time 150 minute sample drawn OGDRAW150M/OGDRAW150S  | : Timer                                  |  |  |  |  |  |  |
| 17. Time 180 minute sample drawn OGDRAW180M / OGDRAW180S (3 hour)   | : Timer                                  |  |  |  |  |  |  |
| 18. The OGTT was (CHECK ONLY ONE): OGCOMPLETE   |  |  |  |  |  |  |  |
| Completed without problem   | 1  |  |  |  |  |  |  |
| Completed <b>with</b> a problem   | 2  |  |  |  |  |  |  |
| Not completed   | 3  |  |  |  |  |  |  |
| If completed without a problem, STOP. Otherwise, complete 19a.  |  |  |  |  |  |  |  |
| <ul> <li>a. Why was the OGTT "completed with a problem" or "not</li> <li>Vomited after glucose load</li> </ul>  | completeds ockromma.                     |  |  |  |  |  |  |
| Fainted or felt ill after glucose load  |  |  |  |  |  |  |  |
| Participant did not drink entire glucose load in 5 minutes  | 3  |  |  |  |  |  |  |
| Difficult blood draw  | 4  |  |  |  |  |  |  |
| Other   | 5  |  |  |  |  |  |  |
| If "Other," Specify:  |  |  |  |  |  |  |  |