



Value

SECTION A: GENERAL STUDY INFORMATION FOR OFFICE USE ONLY

A1. Study ID#: A2. Visit # F/U 12 MonthsV12M

A3. Date Form Completed: ___/___/___
Month Day Year A4. Study Staff Initials: _____

A5. Date of Stress Test: ___/___/___
Month Day Year A6. Stress Tester's Initials: _____

Analysis Variable : rando_dt_TEST_D								
N	N Miss	Mean	SD	Minimum	Lower Quartile	Median	Upper Quartile	Maximum
443	187	-377.2	31.3	-550.0	-391.0	-372.0	-357.0	-291.0

SECTION B: STRESS TEST AND PVR

B1. Voided volume prior to the empty bladder stress test (EBST): _____ mL

Analysis Variable : EBST								
N	N Miss	Mean	SD	Minimum	Lower Quartile	Median	Upper Quartile	Maximum
428	19	178.1	146.3	0.0	60.0	150.0	267.5	790.0

B2. Was the empty bladder stress test positive ...

Yes No

a. ...with Valsalva in dorsal lithotomy position?..... 1 2

EBST_VAL	Frequency	Percent	Cum Freq	Cum Percent
.	6	.	.	.
1:Yes	2	0.45	2	0.45
2:No	439	99.55	441	100.00

Frequency Missing = 6

b. ...with cough in dorsal lithotomy position?

1

2

EBST_COUGH	Frequency	Percent	Cum Freq	Cum Percent
.	5	.	.	.
1:Yes	4	0.90	4	0.90
2:No	438	99.10	442	100.00

Frequency Missing = 5

B3. Record the PVR (volume of _____ mL (Per catheterized measurement; bladder scan may be used if EBST positive/fill not required.)

REMINDER: If EBST is positive, fill is not required and Stress Test is complete. Skip to C1.

B4. Was bladder filled to 300mL or to an MCC <300mL? 300mL.....1

MCC <300mL.....2 → GO TO B4a

BLAD_FILL	Frequency	Percent	Cum Freq	Cum Percent
.	196	.	.	.
1: 300 mL	367	84.56	367	84.56
2: MCC<300 mL	67	15.44	434	100.00

Frequency Missing = 196

B4a. Record MCC: _____ mL

B5. Was the full bladder stress test positive ...

Yes	No	Not Tested
-----	----	------------

i. ...with Valsalva in dorsal lithotomy position?

1

2

3

FBST_VAL	Frequency	Percent	Cum Freq	Cum Percent
.	12	.	.	.
1: Yes	15	3.45	15	3.45
2: No	416	95.63	431	99.08
3: Not Done	4	0.92	435	100.00

Frequency Missing = 12

ii. ...with cough in dorsal lithotomy position?

1

2

3

FBST_COUGH	Frequency	Percent	Cum Freq	Cum Percent
.	12	.	.	.
1: Yes	25	5.75	25	5.75
2: No	402	92.41	427	98.16
3: Not Done	8	1.84	435	100.00

Frequency Missing = 12

iii. ...with Valsalva standing? 1 2 3

FBST_STAND_V	Frequency	Percent	Cum Freq	Cum Percent
.	12	.	.	.
1: Yes	23	5.29	23	5.29
2: No	390	89.66	413	94.94
3: Not Done	22	5.06	435	100.00

Frequency Missing = 12

iv. ...with cough standing? 1 2 3

FBST_STAND_C	Frequency	Percent	Cum Freq	Cum Percent
.	195	.	.	.
1: Yes	28	6.44	28	6.44
2: No	378	86.90	406	93.33
3: Not Done	29	6.67	435	100.00

Frequency Missing = 195

SECTION C: ANTHROPOMETRIC MEASURE

C1. Weight: _____ lbs

Analysis Variable : WEIGHT								
N	N Miss	Mean	SD	Minimum	Lower Quartile	Median	Upper Quartile	Maximum
438	9	170.9	36.5	94.0	143.0	165.0	193.0	308.0